

(2)

Marine Physical Laboratory

AD-A239 192



Fieberling Guyot Studies

Final Report Prepared
for the Office of Naval Research
Department of the Navy
for Grant N00014-89-J-1054
for the Period 10-01-88 - 04-30-91
Principal Investigator: Peter F. Lonsdale

MPL-U-56/91
July 1991

Approved for public release; distribution unlimited.



University of California, San Diego
Scripps Institution of Oceanography

91-07231



91

079

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) MPL-U-56/91			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION University of California, San Diego		6b. OFFICE SYMBOL (If applicable) MPL	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research Department of the Navy		
6c. ADDRESS (City, State, and ZIP Code) Marine Physical Laboratory Scripps Institution of Oceanography San Diego, California 92152			7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable) ONR	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-89-J-1054		
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) FIEBERLING GUYOT STUDIES					
12. PERSONAL AUTHOR(S) Peter F. Lonsdale, Principal Investigator					
13a. TYPE OF REPORT final report		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) July 1991	
15. PAGE COUNT 3					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
			ocean floor geomorphology, deep-sea bedforms, Fieberling Guyot, Seabeam swaths		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The work under this grant concentrated on interpreting ocean-floor geomorphology, especially by fine-scale studies of sites where relief is being actively created or modified, by tectonism and volcanism (e.g. at plate boundaries and young seamounts) or by bottom current activity. An important focus of the latter aspect was to learn how to use deep-sea bedforms (the clearest evidence of geologically effective bottom currents) to infer characteristics of the forcing flow and the benthic boundary layer, as well as for delineating rates and routes of sediment transport.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Peter F. Lonsdale, Principal Investigator			22b. TELEPHONE (Include Area Code) (619) 534-2855		22c. OFFICE SYMBOL MPL

Fieberling Guyot Studies

Final Report Prepared
for Office of Naval Research
Grant N00014-89-J-1054
For the Period 10-01-88 - 04-30-91
Award Amount: \$351,165
Peter F. Lonsdale (Principal Investigator)



Administrative stamp area with a checkmark and handwritten text 'A-1'.

OBJECTIVE

The work under this grant concentrated on interpreting ocean-floor geomorphology, especially by fine-scale studies of sites where relief is being actively created or modified, by tectonism and volcanism (e.g. at plate boundaries and young seamounts) or by bottom current activity. An important focus of the latter aspect was to learn how to use deep-sea bedforms (the clearest evidence of geologically effective bottom currents) to infer characteristics of the forcing flow and the benthic boundary layer, as well as for delineating rates and routes of sediment transport.

BACKGROUND

Fieberling was first described by Carsola and Dietz (1952) as a extinct volcano 500 miles west of San Diego, with a wave-planed summit now 500-700 m below sea-level. Since 1983 several collections of Seabeam swaths across Fieberling and an isotopically dated lava sample (20 Ma), were conducted as part of an ONR-funded study of seamounts off southern California. Many of these seamounts lie in a hotspot chain which has Fieberling as its oldest member, and Guadalupe as the only other volcano which grew above sea-level, though the summits of several others (Hoke, Stoddard, Jasper) were once within a few hundred meters of the sea surface.

In 1986 a 4-day Deep Tow survey of the 70 km² summit plain of Fieberling Guyot was conducted, funded by an ONR 6.2 contract with definition of the near-bottom magnetic field as its primary purpose. Ancillary geologic and biologic observations were made with the sonars and stereo cameras of the Deep Tow vehicle, and near-bottom currents were measured by attaching current meters to the navigational transponder moorings.

APPROACH

The project included a field experiment to monitor the migration of fields of cohesionless sand bedforms (ripples and larger-scale waves) at 600 m depth on the surface of Fieberling Guyot. Bottom currents and benthic boundary layer physics and biology were measured during this experiment by current meters, CTD profiles,

Marine Physical Laboratory

and BASS (Benthic Acoustic Stress Sensor) tripods. The first year of the project was devoted mainly to survey operations that were essential for site selection and problem definition for this project.

First-Year Field Program

A 5-day surface ship survey, navigated by GPS plus Loran C, provided a complete Seabeam bathymetric map of the guyot side-slopes and base, complementing the existing Deep Tow bathymetry of the summit. At the same time as the Seabeam survey, a Sea MARC II survey was made, 3.5 kHz acoustic profiles were taken (mainly to look for evidence of current scour around the base of the volcano and its satellite cones) and the gravity and magnetic field was measured.

Complete high-resolution bathymetry was a prerequisite for many of the studies in the topographic interactions program. The bathymetric data was used in a quantitative comparison of the geomorphology of guyot side-slopes with the slopes of similar but never-emergent volcanoes in the chain which have already been completely surveyed by Seabeam (e.g. Jasper Seamount). During the bathymetric survey amplitude data from the Seabeam system was processed and displayed for quantitative acoustic studies of bottom reverberation and of the Deep Scattering Layer (by Christian de Moustier).

The gravity field was measured with *Washington's* new Bell gravimeter to provide precise definition of the marine geoid, needed for full interpretation of time-varying (oceanographic) information from satellite altimetry. The guyot's gravity field was also used to interpret the density distribution (and hence the internal structure) of the volcano, using inversion techniques. Gravity-derived inferences about the internal structure were valuable because guyot planation had already exposed shallow parts of the interior (down to about 1 km below the former volcano surface), so geologic inference (downward extrapolation of known dike and plug structures) helps interpret the density distribution. Magnetic data was also inverted to define magnetization distributions and hence help interpret the internal structure.

Second-Year Field Program

The MPL Deep Tow system was used for further near-bottom study of the guyot. A survey on the western part of the summit plain concentrated on photographic mapping of current bedforms on previously located sand patches. In addition to a more thorough Deep Tow survey of part of the summit, a sector of the guyot's side slope was acoustically mapped and photographed. This characterized the slope's steepness, small-scale roughness, surface composition (bare rock, cobbles, talus, sand, mud) from the shelf break to the foot of the volcano.

Investigation of ripple and sand wave migration over the summit plain was an integral part of the benthic boundary layer study that was a major component of the ONR Applied Research Initiative. Success of this study was dependent on collaboration with other investigators simultaneously measuring the fluctuations of current and shear stress in the free flow and boundary layer; conversely, appreciation of the bed geometry and mobility is required for interpretation of the boundary layer physics and ecology and distribution of the benthic infauna.